

TURF FERTILIZER + PRODIAMINE 0.29% Details

NY Product No. 60063-41-82757

Type HERBICIDE
 Use NON-CROP, RESIDENTIAL, ORNAMENTAL
 Status REGISTERED
 Restriction Codes None
 Restriction Conditions None
 Toxicity Statement AQUATIC INVERTEBRATES, FISH
 Signal Word CAUTION
 Precautionary Statement DERMAL
 Formulation SOLID
 Renewal Date 08/31/2018
 Initial Registration Date 09/23/2013

Active Ingredients

AI Code	Active Ingredient Name	Percent of AI
110201	Prodiamine	0.29

NYS Labels/Documents

Document Type	Accepted Date	Label/Document No.
PRIMARY LABEL	09/23/2013	536432



TURF FERTILIZER + 0.29% Prodiamine

For selective pre-emergence control of grass and broadleaf weeds in:
 • Established Turf Grasses (excluding golf course putting greens) and lawns
 • Landscape Ornamentals (including established perennials and wildflower plantings)

ACTIVE INGREDIENT:
 Prodiamine, [N,N'-Di-n-propyl-2,4-dinitro-5-(trifluoromethyl)-m-phenylene]diimine 0.29%
 Inert Ingredients 99.71%
TOTAL 100.00%

18-0-3

GUARANTEED FERTILIZER ANALYSIS

TOTAL NITROGEN (N) 18.0% Urea Nitrogen*

SOLUBLE POTASH (K₂O) 3.0%

TOTAL SULFUR (S) 1.4% Free Sulfur (S)

DERIVED FROM: Urea, Polymer-Coated Sulfur-Coated Urea, Murate of Potash.


CHLORINE (Cl) Max. 2.3%

* 4.0% Slowly Available Nitrogen from Polymer-Coated Sulfur-Coated Urea.

ACCEPTED FOR REGISTRATION
 September 23rd, 2013

New York State Department of Environmental Conservation
 Division of Materials Management
 Pesticide Product Registration

CRABGRASS SEED GERMINATION DATES



Approximate Date
 [Symbol] After May 30
 [Symbol] After May 10
 [Symbol] After April 20
 [Symbol] After March 20
 [Symbol] January 1 to March 20

THIS PRODUCT IS NOT INTENDED FOR RETAIL SALE

In the state of Florida, it is recommended that the user of this product follow the Green Industries and Golf Course Best Management Practices found at: http://www.dep.state.fl.us/water/nonpoint/docs/nonpointBMP_Book_Final.pdf and <http://www.dep.state.fl.us/water/nonpoint/docs/nonpointBMP.pdf>

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

See the following section, Precautionary Statements, Hazards to Humans and Domestic Animals, for additional information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
IF SWALLOWED: Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center, or doctor for further treatment advice.
 Have this product container or label with you when calling a poison control center, or doctor, or when going for treatment. You may also call 1-800-858-7378 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is a selective pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in:

- Established turf grasses (excluding golf course putting greens) and lawns
- Landscape ornamentals and wildflower plantings

This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when it is activated by at least 1/2 inch of rainfall, irrigation, or shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown for: (1) Sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds. Do not graze or feed livestock forage cut from areas treated with this product. Do not apply aerially. Do not apply to golf course putting greens. Do not apply this product through any type of irrigation system.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

Barnyardgrass	Kochia
Bluegrass, Annual (<i>Poa annua</i>)	Lambquarter, common
Carpetweed	Lovage
Chickweed, common	Parrotium (Texas, Fall, Browntop)
Chickweed, mouse ear (from seed)	Pigweed
Crabgrass (large smooth)	Purslane, common
Crowfootgrass	Pursley, Florida
Cupress, Woolly	Rescuegrass ³
Foxtails, Annual	Shepherd's Purse ²
Goosegrass ¹	Signalgrass, Broadleaf
Herbit	Speedwell, Perian
Ichgrass	Sprangletop
Johnsongrass (from seed)	Sourgrass, Prostrate
Jungle rice	Witchgrass
Knawweed	Woodruff, Yellow (from seed)

¹In many areas a single application of 0.65 lb. to 1.5 lb. active ingredient per acre (equal to 5.1 to 11.9 lbs./1000 sq. ft. of this product) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 0.65 lb. to 1.0 lb. a.i. per acre (equal to 5.1 to 7.9 lbs./1000 sq. ft. of this product) followed after 60 to 90 days by a second application at doses that would not exceed those given in the Maximum Annual Rate Table. Do not exceed the maximum rate for turf grass species listed in the Maximum Annual Rate Table above.

²Applications for this weed should be made in late summer, fall, or winter prior to germination.

³Suppression only. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

APPLICATION DIRECTIONS

Apply uniformly with suitable, calibrated application equipment.

ESTABLISHED TURF:

This product is a selective pre-emergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the Maximum Annual Rates section of this label. Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

USE PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns: (1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses. (2) Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seeding secondary roots are in the second inch of soil, not thatch plus soil. (3) Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year. (4) Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury. (5) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control. (6) Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table.

Turf Species:	Maximum Application Rate/Calendar Year of turf fertilizer by turf grass species ¹		
	Lbs. Product/A	Lbs. Product/1000 sq. ft.	Lbs. a./A
Creeping Bentgrass	224	5.1	0.85
Creeping Red Fescue	259	5.9	0.76
Buffalograss	345	7.9	1.0
Kentucky Bluegrass			
Perennial Ryegrass	517	11.9	1.5
Bermudagrass ²			
Bahiagrass			
Centipedeagrass			
Seashore Paspalum			
St. Augustinegrass			
Tall Fescue (including turf-type)			
Zoysia			

¹These are the maximum rates per calendar year by species limitations.
²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 0.5 lb. a./acre (equal to 4.0 lbs./1000 sq. ft. of this product). Newly sprigged or plugged Bermudagrass also rooting may be temporarily tolerated. Suppression only of Foxtail, Goosegrass, and Rescuegrass due to reduced product rates used in sprigging situations.

³Do not apply more than 1.5 lbs. a.i. per calendar year per acre (equal to 11.9 lbs./1000 sq. ft. of this product).
 • Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the Maximum Annual Rates Table.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seeding secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the following table below for best overseeding/reseeding results.

Lbs Product/Acre	Lbs Product/1000 sq. ft.	Lbs a./A	Months Before Overseeding		
			North	Transition	South
172	4.0	0.50	4	4	4
224	5.1	0.65	5	4	4
259	5.9	0.75	6	5	5
276	6.3	0.80	-	6	6
345	7.9	1.00	-	7	7
393	9.0	1.14	-	-	8
448	10.3	1.30	-	-	10
517	11.3	1.50	-	-	12

SPREADER SETTINGS

Spreader settings vary by make and model of spreader. It is recommended that individual spreaders are calibrated for the specific product that is to be applied. A walking speed of 3 miles per hour is recommended.

Spreader Model	SPREADER SETTINGS				
	Application Rates ¹		a.i. lbs./Acre (Product lbs./1000 sq. ft.)		
LESCO 021820	0.5	4.0	0.75	5.9	1.0 (7.9)
	D 1/2	E 1/2	G 1/2	1 1/2	
Lely HR	5.25	6	7	8.5	
Vicon 203	15	17	21	27	

¹Application Rates for example only - do not exceed the maximum annual application rate of 1.5 lb. a./Acre (11.9 lb product/1000 sq. ft.)

STORAGE AND DISPOSAL

STORAGE: Store this product in its original container in a dry, cool, secured area. Do not contaminate water, foodstuffs, feed, or seed by storage or disposal.

PRODUCT DISPOSAL: As a responsible environmental practice, where possible, it is recommended that all of the contents of the bag be used, carefully following label directions and precautions.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling, if available, or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned stay out of smoke.

#901833 Net Weight 50 lbs. (22.67 kg)





TURF FERTILIZER + 0.29% Prodiamine

LANDSCAPE ORNAMENTALS

Including Established Perennials and Wildflower Plantings
This product may be applied for residual pre-emergence weed control in ornamentals.

Use rates and When to Apply

Apply in fall and/or spring, 0.65 to 1.5 lbs. active ingredient per acre, which is equal to 5.1 to 11.9 lbs. of this product per 1,000 sq.ft. of ornamental plantings; therefore 20 lbs of this product will treat 1,681 to 3,922 sq.ft. Use the higher rates of application for longer control periods. Sequential applications may be made as long as the total amount of product applied does not exceed the maximum annual application rate of 1.5 lbs. a.i. per acre.

Established Landscape Ornamentals

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate it into the soil.

Newly Transplanted Landscape Ornamentals

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grfts have taken to avoid any inhibition of the tissue union.

Tolerant Ornamental Species

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when it is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

Scientific name	Common name (Tested varieties)	Scientific name	Common name (Tested varieties)	Scientific name	Common name (Tested varieties)	Scientific name	Common name (Tested Varieties)
<i>Abies</i> spp.	Fir	<i>Coloniaestier microphyllus</i>	Rockspray cotoneaster	<i>Juglans</i> spp.*	Walnut*	<i>Pistacia</i> spp.*	Platanus*
<i>Abelia grandiflora</i>	Abelia; Silverwood	<i>Craetagus</i> spp.	Hawthorne	<i>Juniperus brachyopoda</i>	Shrimp plant	<i>Pittosporum rhomboidatum</i>	Queensland pittosporum
<i>Acer palmatum</i>	Japanese maple	<i>Crossosoma</i> spp.	Lucifer	<i>Lagerstroemia indica</i>	Crape myrtle	<i>Pittosporum tobira</i>	Japanese pittosporum
<i>Acer platanoides</i>	Norway maple	<i>Cyperus sanguinolentus</i>	Italian cypress	<i>Lantana maculivariolosa</i>	Weeping lantana	<i>Podocarpus neriifolius</i>	Japanese yew
<i>Achillea</i> sp.	Yarrow (King Edward)	<i>Delosperma alba</i>	White trailing ice plant	<i>Lavender</i> spp.	Lavender (Munstead)	<i>Prunus laurocerasus</i>	English laurel
<i>Actinidia chinensis</i> *	Kiwi*	<i>Delphinium</i> spp.	Larkspur (Blue elf)	<i>Leopodium alpinum</i>	Edelweiss	<i>Prunus</i> spp.*	Almond, Apricot, Nectarine, Peach, Plum, and Prune*
<i>Agapanthus africanus</i>	Lily-of-the-Nile; African Lily	<i>Dianthus barbatus</i>	Maiden pink (Zing)	<i>Ligustrum amurense</i>	Amur privet	<i>Pseudotsuga menziesii</i> **	Douglas-fir**
<i>Akiba quitifolia</i>	Five-leaf or Choochole vine	<i>Dianthus barbatus</i>	Cheddar pink	<i>Ligustrum japonicum</i>	Japanese privet	<i>Pyracantha coccinea</i>	Firethorn, scarlet
<i>Alium caeruleum</i>	Lady's teak or Nodding onion	<i>Dodonaea viscosa</i>	Hop bush	<i>Ligustrum lucidum</i>	Glossy privet (Wax-Leaf)	<i>Pyracantha fortuneana</i>	Firethorn
<i>Anemone hybrida</i>	Japanese anemone	<i>Echinacea purpurea</i>	Coneflower, purple (Magnus)	<i>Ligustrum sinense</i>	Chinese privet (Veriegata)	<i>Pyracantha koidzumii</i>	Firethorn
<i>Aquilegia</i> spp.	Columbine (Red and gold)	<i>Elaeagnus pungens</i>	Silverberry	<i>Lilium</i> spp.	Lily (Lazzi)	<i>Pyrus</i> spp.	Bradford pear
<i>Arctostaphylos densiflora</i>	Vine Hill manzanita	<i>Euonymus fortunei</i>	Wintercreeper	<i>Liriodendron</i>	Liriodendron (Big blue, Light, Silver mound, Evergreen glen)	<i>Quercus rubra</i>	Red oak
<i>Arctostaphylos densiflora</i>	Cape weed	<i>Euonymus japonica</i>	Japanese spindle tree; Evergreen euonymus	<i>Liriodendron var. variegata</i>	Liriodendron, variegated	<i>Quercus shumardii</i>	Shumard oak
<i>Artemisia</i> spp.	Wormwood (Silver mound; Castle)	<i>Euonymus kiautschowica</i>	Spreading euonymus	<i>Liriodendron sinense</i>	Liriodendron, creeping	<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Aster</i> sp. or <i>Symphoricarpos</i> sp.	Aster (Bonny blue, Purple dome)	<i>Fatsia japonica</i>	Japanese aralia	<i>Lobelia cardinalis</i>	Cardinalflower	<i>Rhododendron</i> (including Azalea)	Rhododendron & Azalea (Coral Bells, Delaware valley white, Flame creeper, George L. Tabor, P.J.M., Roseum Elegans, Formosa, Hino-crimson, Wakabitsu, Gumpo)
<i>Aster X frikartii</i>		<i>Forsythia intermedia</i>	Border forsythia	<i>Lonicera japonica</i>	Japanese honeysuckle	<i>Rosa banksiae</i>	Lady Bank's Rose
<i>Athyrium filix-femina</i>	Lady fern	<i>Forsythia suspensa</i>	Weeping forsythia	<i>Lonicera tatarica</i>	Tatarian honeysuckle	<i>Rosmarinus officinalis</i> *	Rosemary*
<i>Aucuba japonica</i>	Japanese Aucuba	<i>Forsythia viridissima</i>	Greenstam forsythia	<i>Laropetalum chinense</i>	Burgundy	<i>Rubus</i> spp.	Black-eyed Susan
<i>Begonia</i> spp.	Fibrous begonia (Hardy Grandis)	<i>Gaillardia</i> spp.	Blanket flower (Goblin)	<i>Lythrum</i> spp.	Loosestrife (Modern Pink)	<i>Rumex crispus</i>	Leathertail Fern
<i>Barberis glandulosa</i>	Barberry	<i>Gardenia jasminoides</i>	Gardenia, Cape-jasmine	<i>Magnolia</i> spp.**	Magnolia*	<i>Santolina viridis</i>	Goldstrum Santolina
<i>Barberis julianae</i>	Winegreen barberry	<i>Gentiana</i>	Gentian	<i>Malosiphia luteola</i>	Ice plant	<i>Saxifraga</i> spp.	Saxifrage, Purple dome
<i>Barberis menziesii</i>	Mentor barberry	<i>Geranium chinense</i>	Cranesbill	<i>Malus</i> spp.*	Crabapple*	<i>Sedum album</i>	Stonecrop
<i>Barberis thunbergii</i>	Japanese barberry	<i>Glaucium</i> spp.**	Gladiolus	<i>Miscanthus sinensis</i>	Yaku ima, Silberfeder	<i>Sedum caeruleum</i>	Stonecrop, Ikedansa
<i>Barberis variegata</i>	Warty barberry	<i>Gynopodium repens</i>	Baby's breath	<i>Nandina domestica</i>	Heavenly bamboo	<i>Sedum spectabile</i>	Stonecrop
<i>Berberis cordata</i>	Red barberry	<i>Hedera helix</i>	English ivy	<i>Narcissus</i> spp.**	Narcissus**	<i>Syringia penicillata</i>	Japanese boxhoney
<i>Boltonia asteroides</i>	Snowbank	<i>Helianthemum</i> spp.	Sunrose	<i>Nerium oleander</i>	Oleander	<i>Taxus cuspidata</i>	Japanese yew
<i>Bougainvillea</i> spp.	Bougainvillea	<i>Humacollis</i> spp.	Daisy (Aztec gold, Stella d'Oro, Tender love)	<i>Olea europaea</i> **	Olive*	<i>Taxus media</i>	Yew
<i>Buddleia davidii</i>	Butterfly-bush (Dwarf Blue, Royal Red)	<i>Hibiscus</i> spp.**	Rose of Sharon**	<i>Ophiopogon japonicus</i> **	Mondo Grass**	<i>Teucrium</i> spp.	Germender
<i>Buxus microphylla</i>	Japanese boxwood	<i>Hibiscus</i> spp.	Meibow (Disco belle white)	<i>Osmantinus heterophyllus</i>	Osmantinus; False holly (Gulf tree)	<i>Thalictrum digitato-carpum</i>	Meadow rue
<i>Callistemon viminalis</i>	Weeping bottlebrush	<i>Hibiscus Rosa-sinensis</i> **	Chinese hibiscus**	<i>Osteospermum frutescens</i>	Trailing African Daisy	<i>Thuja occidentalis</i>	American arborvitae
<i>Calluna vulgaris</i>	Scotch heather	<i>Hosta plantaginea</i>	Hosta, Plantain fly	<i>Oxydendron luteum</i>	Sourwood	<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Campanula carpatica</i>	Tussock bellflower (White Cljas)	<i>H. sieboldiana</i>	Hosta (Seasucker)	<i>Persea americana</i>	Avocado*	<i>Tsuga canadensis</i>	Canada hemlock
<i>Campsis X tagliabuana</i>	Trumpet creeper; Trumpet flower (Madame Galen)	<i>Houttuynia cordata</i> var. <i>variegata</i>	Bigleaf hydrangea	<i>Phloxia traseri</i>	Fraser's phlox; Redtop	<i>Tulipa</i> spp.	Tulip
<i>Carpobrotus edulis</i>	Hottentot fig; Ice plant	<i>Hydrangea macrophylla</i>	Bigleaf hydrangea	<i>Pitysostegia virginiana</i>	False dogonhead (Wick)	<i>Viburnum japonicum</i>	Japanese viburnum
<i>Cassia artemisioides</i>	Foothill Cassia	<i>Ilex cornuta</i> **	Chinese holly**	<i>Picea</i> spp.**	Spruce species** (Colorado blue, Norway, etc.)	<i>Viburnum odoratissimum</i>	Sweet viburnum
<i>Ceanothus rigidus</i>	Wild Lilac	<i>Ilex crenata</i>	Japanese holly	<i>Pieris japonica</i>	Lily-of-the-valley shrub	<i>Viburnum plicatum</i>	Japanese snowball
<i>Cercalostigma plumbaginoides</i>		<i>Ilex opaca</i>	American holly	<i>Pinus brutia</i>	Calabrian pine	<i>Viburnum rigidum</i>	Canary Island Viburnum
<i>Chamaecyparis pisifera</i>	False Cypress	<i>Ilex parvifolia</i>	Holly	<i>Pinus canadensis</i>	Canary Island pine	<i>Viburnum tinus</i>	Laurustinus
<i>Chrysanthemum nipponicum</i>	Chrysanthemum	<i>Ilex vomitoria</i>	Yaupon holly	<i>Pinus elliotii</i>	Slash pine	<i>Viburnum trilobum</i>	Cranberry Bush
<i>Cleyera japonica</i>	Cleyera	<i>Inula ensifolia</i>		<i>Pinus halepensis</i>	Aleppo pine	<i>Viburnum wrightii</i>	Leatherleaf Viburnum
<i>Citrus</i> spp.*	Citrus species*	<i>Iris ensata</i>	Sword-leaved iris (Jodestong)	<i>Pinus nigra</i>	Austrian black pine	<i>Vinca major</i>	Greater periwinkle, Vinca
<i>Carex</i> spp.	Carex; Calliopsis (Early sunrise, Moonbeam)	<i>Iris sibirica</i>	Siberian iris (Cabernet)	<i>Pinus palustris</i>	Longleaf pine	<i>Vinca minor</i>	Common periwinkle
<i>Cornus florida</i>	Flowering dogwood	<i>Iris</i> spp.	Iris**	<i>Pinus radiata</i>	Menturey pine	<i>Vitis</i> spp.*	Grape*
<i>Cornus sibirica</i>	American dogwood	<i>Jasminum nudiflorum</i>	Winter jasmine	<i>Pinus strobus</i>	Eastern white pine	<i>Weigela florida</i>	Old fashioned Weigela
<i>Coriaria selkoeana</i>	Pampas grass	<i>Juniperus chinensis</i>	Chinese juniper (Nick's conical)	<i>Pinus sylvestris</i>	Scotch pine	<i>Yucca aloifolia</i>	Spanish bayonet
<i>Cotoneaster argenteus</i>	Cranberry cotoneaster	<i>Juniperus conferta</i>	Shore juniper (Blue Pacific)	<i>Pinus taeda</i>	Loblolly pine	<i>Yucca filamentosa</i>	Yucca, Adam's needle
<i>Cotoneaster boxifolius</i>	Cotoneaster	<i>Juniperus davurica</i>	Juniper	<i>Pinus thunbergiana</i>	Japanese black pine		* Do not use on food producing trees, vines, or plants.
<i>Cotoneaster damieri</i>	Bearberry cotoneaster	<i>Juniperus horizontalis</i>	Creeping juniper	<i>Pinus virginiana</i>	Virginia pine		** Not for use on conifer grown plants

CONDITIONS OF SALE AND WARRANTY

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Turf Care Supply Corp. or the Seller. All such risks shall be assumed by the Buyer. Turf Care Supply Corp. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. **Turf Care Supply Corp. makes no other express or implied Warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Turf Care Supply Corp. or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this Product.** Turf Care Supply Corp. and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of Turf Care Supply Corp.

<p>EPA REG. NO. 60063-41-82757 EPA EST. NO. 82757-OH-001 (Lot # Begins with MF) EPA EST. NO. 82757-FL-001 (Lot # Begins with S) EPA EST. NO. 82757-MA-002 (Lot # Begins with H)</p> <p>#901833</p>	<p>F1612</p> <p>Distributed by: Turf Care Supply Corp. 50 Pearl Road, Suite 200 Brunswick, OH 44212</p>	
---	---	--